

Seasonal Patterns in Aquatic Bird Counts at Five Andean Lakes of Ecuador

Author(s): Esteban A. Guevara , Tatiana Santander and Joost F. Duivenvoorden

Source: *Waterbirds*, 35(4):636-641. 2012.

Published By: The Waterbird Society

URL: <http://www.bioone.org/doi/full/10.1675/063.035.0413>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

Seasonal Patterns in Aquatic Bird Counts at Five Andean Lakes of Ecuador

ESTEBAN A. GUEVARA^{1,2,*}, TATIANA SANTANDER¹ AND JOOST F. DUIVENVOORDEN²

¹Aves&Conservación – BirdLife in Ecuador, Pasaje Joaquín Tinajero E3-05 and Jorge Drom, Casilla 17-17 906, Quito, Ecuador

²Institute for Biodiversity and Ecosystem Dynamics, Universiteit van Amsterdam, Science Park, 1098 HX, Amsterdam, The Netherlands

*Corresponding author; E-mail: esalguan@yahoo.com

Abstract.—Seasonal (semestral) counts of aquatic birds at five Andean lakes in Ecuador (Colta, La Mica, Yambo, Yahuarcocha and San Pablo) were analyzed to detect trends in population size between 2004 and 2011. Trends of four abundant species (*Ardea alba*, *Anas georgica*, *Fulica ardesiaca* and *Oxyura jamaicensis*) and those of five functional groups (based upon body size and diet) were tested using generalized additive mixed modeling. In total, 19 bird species were consistently recorded over the whole period. The Andean Coot (*F. ardesiaca*) was found in highest densities (c. 3 birds per ha at Colta and Yambo). No significant trends were detected. However, the counts of functional groups exhibited a seasonal pattern (higher counts in February than in July), possibly due to lower resource availability in the dry season. Piscivorous species were relatively abundant at one lake (Yahuarcocha), which might be related to the introduction of Tilapiine cichlid fishes. Received 29 February 2012, accepted 26 June 2012.

Key words.—Andes, bird monitoring, functional groups, GAMM.

Waterbirds 35(4): 636-641, 2012

The inter-Andean valleys of Ecuador encompass wetlands that sustain important populations of aquatic bird species (Ridgely and Greenfield 2001). In Ecuador, a monitoring program to provide baseline information on the distribution and abundance of aquatic birds started in 2004 (López-Lanús and Blanco 2005). Such data are essential to provide guidelines for the management and conservation of wetlands (Tamisier and Boudouresque 1994; Paillisson *et al.* 2002; Lagos *et al.* 2008; Mistry *et al.* 2008). Andean lakes have endemic fish (Ruzzante *et al.* 1998) and bird species that are threatened at the national level (e.g. *Theristicus melanopis* Williams and Olmedo 2002). Bird monitoring data for these lakes are especially important because the increasing extraction of water for drinking and irrigation (Santander *et al.* 2006) and alterations in trophic cascades due to the introduction of exotic fish, may precipitate a decline in bird populations (Ortubay *et al.* 2006). In this context, the aim of our study was to assess trends in the first eight years of seasonal (semestral) bird counts for five inter-Andean lakes of Ecuador.

METHODS

Study Area

Lake Colta is located 17 km south of the city of Riobamba (Chimborazo province, 1°44'S 78°45'W) at 3,420 m above sea level (a.s.l.). Lake Colta covers 240 ha and has a maximum depth of 3.5 m (Santander *et al.* 2006). La Mica is a 360 ha glacial lake with an average depth of 22.5 m situated in the Antisana Ecological Reserve (Napó province, 00°32'S 78°12'W) at 3,900 m a.s.l. Lake Yahuarcocha is in the dry inter-Andean valley of northern Ecuador (Imbabura province, 00°22'N 78°06'W) at 2,210 m a.s.l. The lake covers 230 ha at a maximum depth of 7.9 m (Consejo Provincial de Imbabura 1997). Lake San Pablo is 20 km south of Lake Yahuarcocha (Imbabura province, 0°13' N, 78°14' W) at 2660 m a.s.l. With an area of 670 ha and a maximum depth of 35 m, it is one of the largest Andean lakes in Ecuador (Santander *et al.* 2006). Lake Yambo is situated in the inter-Andean valley of Central Ecuador (Cotopaxi province, 1°05'S 78°35'W), at 2,600 m a.s.l. The lake covers 16 ha and has a maximum depth of 25 m (Steinitz-Kannan *et al.* 1993).

Bird Census

At each lake, bird counts were performed in February and July during 2004-2011. On a typical monitoring day, experienced observers carried out their census, which started in the morning (08:00 h) and lasted for two to three hours. Observations were conducted from either a boat that toured the wetlands perimeter at a constant velocity, remaining at 60-100 m from the shore

to avoid massive take-off of birds that congregate, or by hiking along the lake shores. If more than one observer counted birds simultaneously the highest value was considered.

Numerical Analysis

Because the total monitoring period was short, we included the lakes as random effects and analyze the counts of only the four most abundant species (*Ardea alba*, *Anas georgica*, *Fulica ardesiaca* and *Oxyura jamaicensis*). These four species are common in Andean lakes (Fjeldså and Krabbe 1990). Also, we tested for trends in the counts of all species grouped into five functional groups related to body size and diet (large and medium size piscivorous and medium size herbivorous, insectivorous and omnivorous; del Hoyo *et al.* 1992, 1996).

Most count-time relationships were distinctly non-linear and the variances in the counts differed substantially between species or functional groups (Fig. 1a, 1b). Therefore, we applied a generalized additive mixed model of counts against time, applying one model for the counts of the selected species, and another model for the counts of the functional groups (Fewster *et al.* 2000; Zuur *et al.* 2009). The starting models were as follows:

```
model <- gamm(sqrt(counts) ~ ID + cos(time*pi)
+ s(time, by = ID)), method = "ML", random =
list(IDlake=~1), weights = varIdent(form = ~ 1 | ID))
```

In this, ID refers to the selected species or functional groups as factor; $\cos(\text{time} \cdot \pi)$ a cosinus function to implement a fixed seasonality effect in the model; $s(\text{time}, \text{by} = \text{ID})$ a thin plate regression spline smoother, one for each level of ID; $\text{random} = \text{list}(\text{IDlake} \sim 1)$ the name of each lake as random effect; $\text{method} = \text{"ML"}$ the Maximum Likelihood method of model fitting; and $\text{weights} = \text{varIdent}(\text{form} = \sim 1 | \text{ID})$ to the implementation of different variances for each level of ID in the model. We applied the square root transformation of the counts because the weights argument in the nlme function (which forms part of the gamm function) may only be used in case of a gaussian identity link. The value of Akaike's Information Criterion (AIC) of the above model was similar (selected species) to or lower (functional groups) than models from which the seasonality effect was removed. The standardized residuals of both models (selected species and functional groups) showed clear patterns of short-term autocorrelation. Therefore, we adjusted the models by including the following term for the correlation structure: $\text{corARMA}(p = 3, q = 3)$. As a result, both models showed substantial lower AIC values, and the autocorrelation of residuals was eliminated (functional groups) or highly reduced (selected species). Further model validation (residuals against fitted values or against time) did not produce anymore pattern. The residuals did not deviate strongly from normality (visual inspection of quantile-quantile plots per lake). Analyses were done using R 2.14.1 (R Development Core Team 2011), applying the mgcv (Wood 2006), nlme (Pinheiro *et al.* 2011) and lattice (Sarkar 2008) packages.

RESULTS

A total of 19 bird species were recorded and classified into five functional groups (Table 1). Mean (\pm one SD) seasonal bird counts were 963 ± 388.1 at La Mica, 919 ± 229.6 at San Pablo, 831 ± 327.4 at Colta, 797 ± 277.2 at Yahuarcocha and 310 ± 141.7 at Yambo. Mean (\pm SD) seasonal counts per ha were 19.4 ± 8.9 at Yambo, 3.5 ± 1.2 at Yahuarcocha, 3.5 ± 1.4 at Colta, 2.7 ± 1.1 at La Mica and 1.5 ± 0.4 at San Pablo. *F. ardesiaca* exhibited high densities, especially at Yambo and Colta (approximately 3 birds per hectare at both lakes).

Between 2004 and 2011, no significant trends were detected in the counts of the selected species or functional groups (Table 2). The seasonality effect (cosinus function) was highly significant for functional groups ($P < 0.001$) reflecting consistently higher counts in February than in July (Table 2). By including lakes as random effects, the trends in selected species or functional groups in individual lakes were not tested. Yet, the counts of *F. ardesiaca* at the lakes San Pablo, La Mica and Colta, and those of the large size piscivorous group at Lake Yahuarcocha and the medium size omnivorous group at lakes San Pablo and La Mica showed a distinct upward tendency, whereas those of *A. georgica* at Lake La Mica, *O. jamaicensis* in Lake San Pablo and the herbivorous group at Lake Yambo seemed decreasing in time.

DISCUSSION

The high densities of *F. ardesiaca* are consistent with other studies that report species of the *Fulica* genus as most abundant in wetlands (Hamdi *et al.* 2008). The tendency for large piscivorous birds to increase at Yahuarcocha is noteworthy, also piscivorous species seem more abundant at this lake compared to the other lakes (c. one bird per ha at Yahuarcocha versus 0.0 to 0.04 birds per ha at the other lakes). Possibly, these results may relate to the recent introduction of Tilapine cichlid fish (*Oreochromis niloticus* and *Tilapia mossambica*) at Yahuarcocha. The introduction of these fish has been postulated to explain the presence and growth of




Figure 1. Seasonal (semestral) bird counts in five Andean lakes in Ecuador between 2004 and 2011, for each of the selected species (Fig. 1a) and functional groups (Fig. 1b). The lines give the loess smoother (applying a span of 0.9).

Table 2. Results of Generalized Additive Mixed Modeling analysis for selected species and functional groups.

Selected species			
Parametric Terms:			
	df	F	p-value
cos(time * pi)	1	3.24	0.073
ID (selected species)	3	25.6	1.5E-14
Approximate significance of smooth terms			
	Estimated df	F	p-value
s(time): <i>A. alba</i>	0.9978	0.33	0.57
s(time): <i>A. georgica</i>	0.9990	0.022	0.88
s(time): <i>F. ardesiaca</i>	0.9999	0.23	0.63
s(time): <i>O. jamaicensis</i>	0.9999	0.017	0.90
Functional groups			
Parametric Terms:			
	df	F	p-value
cos(time * pi)	1	17.8	3.1E-05
ID (functional group)	4	33.7	<2e-16
Approximate significance of smooth terms			
	Estimated df	F	p-value
s(time): Piscivorous, large	1	0.96	0.33
s(time): Piscivorous, medium	1	0.68	0.41
s(time): Herbivorous	1	0.029	0.87
s(time): Insectivorous	1	0.017	0.90
s(time): Omnivorous	1	0.053	0.82

Phalacrocorax brasilianus and other piscivorous birds from coastal origin at Lake Yahuarcocha (Jahn *et al.* 2010; Guevara *et al.* 2011; Santander *et al.* 2011). Recent observations of aquatic bird species outside their formerly known range have been suggested as an indication of the ongoing colonization of Andean wetlands (Henry 2012). Given the distinct upward or downward tendencies in the counts of several species or functional groups at a number of lakes, continued monitoring should enable researchers to provide firmer evidence for changing waterbird population sizes.

Seasonality patterns in neotropical bird counts have mostly been reported for terrestrial bird communities (e.g. Karr 1976; Loiselle and Blake 1992; see also Alves and Pereira 1998). Seasonal variations in rainfall and temperature have been identified as primary factors driving food resource availability in tropical systems (e.g. seeds, fruits, insects; Karr 1976; Faaborg *et al.* 1984), which ultimately influence bird abundances. Inter-

polated climate surfaces developed for the period between 1950 and 2000 (Hijmans *et al.* 2005) indicate that the precipitation in July and August at the five lakes analyzed in this study is relatively low. Therefore, a hypothetical decline in resource abundance during the dry season might explain the seasonality pattern of the species clustered in functional groups.

ACKNOWLEDGMENTS

We thank Wetlands International and its Neotropical Waterbird Census Programme, and the Government of Imbabura Province for supporting our fieldwork. We are grateful to the staff of Aves&Conservación and the volunteers who participated in the censuses. The present paper is part of EAG's M.Sc. study in Tropical Ecology at the University of Amsterdam, which was funded by the National Secretary of Science, Technology and Higher Education of Ecuador SENESCYT.

LITERATURE CITED

Alves, M. A. S. and E. F. Pereira. 1998. Richness, abundance and seasonality of bird species in a lagoon of

- an urban area (Lagoa Rodrigo de Freitas) of Rio de Janeiro, Brazil. *Ararajuba* 6: 110-116.
- Consejo Provincial de Imbabura. 1997. Manejo sustentable de la laguna de Yahuarcocha. Consejo Provincial de Imbabura, Ibarra, Ecuador.
- del Hoyo, J., A. Elliott and J. Sargatal. 1992. Handbook of the birds of the world, Vol. 1: Ostrich to Ducks. Lynx Edicions, Barcelona, Spain.
- del Hoyo, J., A. Elliott and J. Sargatal. 1996. Handbook of the birds of the world, Vol. 3: Hoatzin to Auks. Lynx Edicions, Barcelona, Spain.
- Faaborg, J., W. J. Arendt, and M. S. Kaiser. 1984. Rain-fall correlates of bird population fluctuations in a Puerto Rican dry forest: A nine year study. *Wilson Bulletin* 96: 575-593.
- Fewster, R. M., S. T. Buckland, G. M. Siriwardena, S. R. Baillie and J. D. Wilson. 2000. Analysis of population trends for farmland birds using generalized additive models. *Ecology* 81: 1970-1984.
- Fjeldså, J. and N. Krabbe. 1990. Birds of the High Andes. Zoological Museum, University of Copenhagen and Apollo Books, Svendborg, Denmark.
- Guevara, E. A., T. Santander, T. Mueces, K. Teran and P. Henry. 2011. Population growth and seasonal abundance of the Neotropical Cormorant (*Phalacrocorax brasilianus*) at highland lakes in Ecuador. *Waterbirds* 34: 499-503.
- Hamdi, N., F. Charfi and A. Moali. 2008. Variation of the waterbird community relying to the Ichkeul National Park, Tunisia. *European Journal of Wildlife Research* 54: 417-424.
- Henry, P. Y. 2012. Distributional and altitudinal range extensions for birds from Ecuador. *Boletín SAO* 20: 89-106.
- Hijmans, R. J., S. E. Cameron, J. L. Parra, P. G. Jones and A. Jarvis. 2005. Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* 25: 1965-1978.
- Jahn, O., P. Cosgrove, C. Cosgrove, T. M. Cevallos and T. Santander. 2010. First record of Brown Pelican (*Pelecanus occidentalis*) from the Ecuadorian highlands. *Cotinga* 32: 108.
- Karr, J. R. 1976. Seasonality, resource availability, and community diversity in tropical bird communities. *American Naturalist* 110: 973-994.
- Lagos, N. A., P. Paolini, E. Jaramillo, C. Lovengreen, C. Duarte and H. Contreras. 2008. Environmental processes, water quality degradation and decline of waterbirds populations in the Rio Cruces wetland, Chile. *Wetlands* 28: 938-950.
- Loiselle, B. A. and J. G. Blake. 1992. Population variation in a tropical bird community. *BioScience* 42: 838-845.
- López-Lanús, B. and D. Blanco. 2005. El censo neotropical de aves acuáticas 2004. Global Series N° 17. Wetlands International. Buenos Aires, Argentina.
- Mistry, J., A. Berardi and M. Simpson. 2008. Birds as indicators of wetland status and change in the North Rupununi, Guyana. *Biodiversity Conservation* 17: 2383-2409.
- Ortubay, S., V. Cussac, M. Battini, J. Barriga, J. Aigo, M. Alonso, P. Macchi, M. Reissig, J. Yoshioka and S. Fox. 2006. Is the decline of birds and amphibians in a steppe lake of northern Patagonia a consequence of limnological changes following fish introduction? *Aquatic Conservation* 16: 93-105.
- Paillisson, J. M., S. Reeber and L. Marion. 2002. Bird assemblages as bio-indicators of water regime management and disturbance in natural wet grasslands. *Biological Conservation* 106: 115-127.
- Pinheiro, J., D. Bates, S. DebRoy, D. Sarkar and the R Development Core Team. 2011. nlme: Linear and nonlinear mixed effects models. R package version 3.1-103.
- R Development Core Team. 2011. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing, Vienna, Austria.
- Ridgely, R. S. and P. Greenfield. 2001. The Birds of Ecuador. Cornell University Press, Ithaca, New York.
- Ruzzante, D. E., S. J. Walde, V. E. Cussac, P. J. Macchi and M. F. Alonso. 1998. Trophic polymorphism, habitat and diet segregation in *Percichthys trucha* (Pisces: Percichthyidae) in the Andes. *Biological Journal of the Linnean Society* 65: 191-214.
- Santander, T., B. Haase and J. R. Hidalgo. 2006. Advancing a range-wide approach to waterbird conservation at priority sites throughout the Neotropics, Capítulo Ecuador. Report to BirdLife International and U. S. Fish & Wildlife Service. Aves&Conservación, Quito, Ecuador.
- Santander, T., K. Terán, T. Mueces, A. Lara, C. Llumiquire and E. Guevara. 2011. Registros inusuales de aves costeras en lagunas Altoandinas de Ecuador. *Cotinga* 33: 105-107.
- Sarkar, D. 2008. Lattice: Multivariate data visualization with R. Springer, New York, New York.
- Steinitz-Kannan, M., M. Nienber, M. Riedinger and R. Kannan. 1993. The fossil diatoms of Lake Yambo, Ecuador. A possible record of el Niño events. *Bulletin de l'Institut Français d'Etudes Andines* 22: 227-241.
- Tamisier, A. and C. Boudouresque. 1994. Aquatic bird populations as possible indicators of seasonal nutrient flow at Ichkeul Lake, Tunisia. *Hydrobiologia* 280: 149-156.
- Williams, R. and I. Olmedo. 2002. Bandurria Carinegra (Theristicus melanopus). Pages 72-73 in Libro rojo de las aves del Ecuador (T. Granizo, C. Pacheco, M. B. Ribadeneira, M. Guerrero and L. Suárez, Eds.). SIMBIOE, Conservacion Internacional, EcoCiencia, Ministerio del Ambiente, UICN Serie de Libros Rojos del Ecuador tomo 2, Quito, Ecuador.
- Wood, S. N. 2006. Generalized additive models: An introduction with R. Chapman and Hall/CRC Boca Raton, Florida.
- Zuur, A. F., E. Ieno, N. J. Walker, A. A. Savaliev and G. M. Smith. 2009. Mixed effects models and extensions in ecology with R. Springer, New York, New York.